



# R2C2139



Relative Maturity: **2.10**  
Genetic Placement: **DEF/off**

Primary Adaptation: **MN, WI**  
Secondary Adaptation: **SD, MI**


- > GENRR2Y with SCN resistance
- > Very good IDC tolerance
- > Rps1c gene for PRR resistance
- > Moderate BSR resistance

## Characteristics

Canopy Type	Int	Flower Color	P	Height	M
Hilum Color	IB	Oil Content	N/A	Pod Color	TN
Protein Content	N/A	PRR Gene	Rps1c	Pubescence Type	GR
SCN Resistance	R3, MR14				

The data presented herein represents the most current information available, but results may vary due to environmental conditions.

© 2008 Winfield Solutions, LLC. All rights reserved. CROPLAN GENETICS is a registered trademark of Land O'Lakes, Inc.



SOYBEANS




# R2C2139



Relative Maturity: **2.10**  
Genetic Placement: **DEF/off**

Primary Adaptation: **MN, WI**  
Secondary Adaptation: **SD, MI**

## Agronomic Data



The data presented herein represents the most current information available, but results may vary due to environmental conditions.

© 2008 Winfield Solutions, LLC. All rights reserved. CROPLAN GENETICS is a registered trademark of Land O'Lakes, Inc.



# R2C2139



Relative Maturity: **2.10**  
Genetic Placement: **DEF/off**

Primary Adaptation: **MN, WI**  
Secondary Adaptation: **SD, MI**

**Rating Scale:**

- 1 = Excellent
- 2-3 = Very Good
- 5 = Fair
- 6-7 = Poor
- 8-9 = Very Poor
- N/A = Not Available

**Genetic Origin:**

E = Eastern, W = Western, N = Northern, S = Southern  
*Capitalization shows dominance*

**SCN Resistance:**

R = Resistant to races listed, MR = Moderately resistant to races listed

**Emergence:**

E = Excellent, VG = Very Good, G = Good

**Canopy Type:**

Nar = Narrow, Int = Intermediate, Bush = Bushy

**Flower Color:**

P = Purple, W = White

**Pubescence Type:**

GR = Gray, TW = Tawny, LTW = Light Tawny

**Pod Color:**

TN = Tan, BR = Brown

**Hilum Color:**

- YE = Yellow/Clear
- GR = Gray
- BL = Black
- IB = Imperfect Black
- BR = Brown
- BF = Buff
- SL = Slate
- TN = Tan

*Protein and oil data collected from three or more location observations in 2006-2007, adjusted to 13 percent moisture.*

*These ratings reflect trends observed in research trials that change with variations in rainfall, temperature, crop production patterns and other factors. Ratings on new varieties are based on limited data and may change as more data is collected.*